

TABLE V
REPORTED DISEASES BY AGE GROUP
CASES PER 100,000 POPULATION¹
2008

	Population	<1	1-4	5-9	10-14	15-19	20-29	30-39	40-49	50-59	60+	TOTAL
AMEBIASIS	397,130	0.8	0.7	1.8	2.2	1.8	1.6	2.0	1.6	0.8	0.3	1.4
ANAPLASMA PHAGOCYTOPHILUM	1,530,851	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
BOTULISM, INFANT ²	1,746,831	2.0										2.0
BOTULISM, WOUND	1,709,339	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
BOTULISM, OTHER	1,866,100	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
BRUCELLOSIS	3,719,315	0.0	0.1	0.1	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0
CAMPYLOBACTERIOSIS	3,594,826	24.7	17.3	8.5	4.8	4.1	4.5	3.6	4.1	5.6	4.8	5.9
CHICKENPOX (VARICELLA)	3,429,814	66.0	61.1	234.8	113.0	8.5	4.4	3.3	2.0	1.2	1.4	32.2
CHOLERA	2,889,253	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
CREUTZFELDT-JAKOB DISEASE	3,443,515	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.3	0.1
CRYPTOSPORIDIOSIS	24,326,974	14.6	56.1	40.7	22.3	5.7	7.9	13.5	6.6	2.5	2.2	13.7
CYCLOSPORIASIS		0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.0
CYSTICERCOSIS		0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0
DENGUE FEVER		0.0	0.0	0.0	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1
EHRLICHIA CHAFFEENSIS		0.0	0.4	0.5	0.1	0.1	0.1	0.0	0.1	0.0	0.1	0.1
ENCEPHALITIS, NONARBOVIRAL		0.0	0.1	0.0	0.0	0.1	0.0	0.1	0.1	0.0	0.2	0.1
ESCHERICHIA COLI, SHIGA TOXIN-PRODUCING (STEC)		11.3	7.5	1.9	1.3	1.1	0.4	0.4	0.5	0.7	0.8	1.4
HAEMOPHILUS INFLUENZAE TYPE B, INVASIVE		0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0
HANTAVIRUS INFECTION		0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
HEMOLYTIC UREMIC SYNDROME		0.3	0.6	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0
HEPATITIS A, ACUTE		0.5	0.5	0.2	0.8	1.4	0.9	0.9	1.5	1.0	1.7	1.1
HEPATITIS B, ACUTE		0.3	0.0	0.1	0.1	0.4	2.3	3.6	4.3	3.6	2.2	2.3
HEPATITIS B, PERINATAL ³		0.8	1.3									1.0
HEPATITIS C, ACUTE		0.0	0.0	0.1	0.0	0.1	0.3	0.5	0.6	0.2	0.0	0.2
HEPATITIS D, ACUTE		0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
INFLUENZA-ASSOCIATED PEDIATRIC MORTALITY ⁴		0.8	0.1	0.1	0.1	0.0						0.1
INFLUENZA, NOVEL A		0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0
LEGIONELLOSIS		0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.3	0.7	1.3	0.3
LISTERIOSIS		1.8	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.2	0.5	0.2
LYME DISEASE		0.0	0.1	0.3	0.7	0.3	0.6	0.8	0.9	0.9	0.5	0.6
MALARIA		0.0	0.3	0.2	0.4	0.4	0.5	0.4	0.3	0.4	0.3	0.4
MENINGITIS, ASEPTIC		125.7	5.5	7.7	7.7	5.4	6.6	5.6	4.7	2.9	3.1	7.2
MENINGITIS, BACTERIAL		21.2	1.7	0.8	0.4	0.7	0.9	1.2	1.8	1.6	2.1	1.6
MENINGITIS, OTHER		0.5	0.0	0.0	0.0	0.1	0.3	0.8	1.0	0.6	0.4	0.4
MENINGOCOCCAL INFECTION		2.3	0.7	0.1	0.1	0.3	0.3	0.1	0.1	0.1	0.5	0.3
MUMPS		0.0	0.2	0.4	0.2	0.1	0.1	0.1	0.0	0.0	0.0	0.1
PERTUSSIS		115.3	15.3	29.8	22.0	4.2	2.2	3.1	2.7	1.8	1.2	8.4
PRIMARY AMOEBOIC MENINGOENCEPHALITIS		0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Q FEVER, ACUTE		0.0	0.1	0.0	0.0	0.1	0.0	0.1	0.1	0.0	0.2	0.1
Q FEVER, CHRONIC		0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0
SALMONELLOSIS		210.5	72.0	29.0	16.3	12.9	14.3	12.2	12.1	15.4	22.4	22.9
SHIGELLOSIS		23.2	98.6	94.9	21.1	5.5	8.3	7.2	4.5	3.6	3.0	19.2
SPOTTED FEVER GP RICKETTSIOSSES		0.0	0.1	0.5	0.3	0.3	0.2	0.3	0.3	0.2	0.2	0.3
STREPTOCOCCUS, GROUP A		4.8	1.8	1.2	0.5	0.2	1.0	0.9	1.5	2.3	4.6	1.8
STREPTOCOCCUS, GROUP B		30.5	0.3	0.1	0.0	0.3	0.7	0.9	1.9	3.8	6.3	2.4
STREPTOCOCCUS PNEUMONIAE		21.7	10.1	3.1	1.2	1.3	1.9	4.1	7.3	12.7	20.6	7.8
TETANUS		0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0
TYPHOID FEVER		0.0	0.3	0.2	0.2	0.1	0.2	0.1	0.0	0.1	0.0	0.1
TYPHUS, MURINE		0.0	0.0	1.2	1.2	0.8	0.3	0.5	0.6	0.8	0.7	0.6
VIBRIO PARAHAEMLYticUS		0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.0	0.1	0.1	0.0
VIBRIO VULNIFICUS		0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.2	0.2	0.1
VIBRIO, OTHER/UNSPECIFIED		0.0	0.0	0.3	0.2	0.1	0.0	0.1	0.0	0.2	0.2	0.1
VISA		0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
WEST NILE FEVER		0.0	0.0	0.1	0.1	0.0	0.1	0.1	0.1	0.2	0.2	0.1
WEST NILE NEUROINVASIVE DISEASE		0.8	0.0	0.0	0.0	0.1	0.1	0.1	0.0	0.2	0.5	0.2
YERSINIOSIS		1.0	0.1	0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.1	0.1

¹ DSHS Center for Health Statistics estimated 2008 Texas population data <http://www.dshs.state.tx.us/chs/popdat/detailX.shtml>.

² Infant botulism rates are calculated using the population under 1 year of age.

³ Perinatal hepatitis B cases are defined as infants >1 months through 24 months of age. The statewide rate was calculated using the population under 2 years of age, which approximates this cohort. The rate calculation for those <1 year does not exclude those <1 month of age and the rate calculation for the 1-4 year cell includes only the 1 year old population.

⁴ Pediatric influenza deaths are defined as deaths occurring in persons under 18 years of age. Rates are calculated for the populations <18 years of age; the rate given in the 15-19 year cell is calculated for the 15-17 year-old population.